

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : A61K 45/05, A01N 63/00, C12N 5/00	A1	(11) International Publication Number: WO 96/16674 (43) International Publication Date: 6 June 1996 (06.06.96)
(21) International Application Number: PCT/US95/15307 (22) International Filing Date: 27 November 1995 (27.11.95) (30) Priority Data: 08/348,659 1 December 1994 (01.12.94) US 08/475,679 7 June 1995 (07.06.95) US (71) Applicant: NEW ENGLAND DEACONESS HOSPITAL CORPORATION [US/US]; One Deaconess Road, CR13, Boston, MA 02115 (US). (72) Inventors: SCADDEN, David, T.; 62 Lexington Street, Weston, MA 02193 (US). FREEDMAN, Andrew; 4 Wynnstay Close, Grangetown, Cardiff 2F1 7NB (GB). (74) Agents: LOREN, Ralph, A. et al.; Lahive & Cockfield, 60 State Street, Boston, MA 02109 (US).	(81) Designated States: CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(54) Title: IN VITRO T-LYMPHOPOIESIS SYSTEM (57) Abstract The present invention provides an <i>in vitro</i> T-lymphopoiesis system in which a population of T-cells is produced from precursor cells expressing CD34. The T-lymphopoiesis system of the present invention produces a population of T-cells of which approximately 17-74 % express CD2, approximately 1.5-34 % express CD3, and approximately 16-61 % express CD4, and approximately 0-15 % express CD8. A method of producing such a population of T-cells <i>in vitro</i> , as well as various compositions including T-cells of the present invention, are also provided.		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

IN VITRO T-LYMPHOPOIESIS SYSTEM

Government Support

The work described herein was supported, in part, by a grant from the National
5 Institutes of Health (HL 44851). The United States government may have certain rights in
the invention.

Background of the Invention

T-lymphocytes, or T-cells, are cornerstones of the mammalian immune system,
10 responsible for cell-mediated immunity against foreign antigens. T-cells, like other
lymphocytes, develop from pluripotent hematopoietic stem cells produced primarily in
hematopoietic tissues (i.e. in the liver in fetuses and in bone marrow in adults). Some of
these precursor stem cells migrate through the blood to the thymus, and T-cell differentiation
occurs there.

15 The majority of T-lymphocytes are immune system regulators, known as helper T
cells and suppressor T cells, that act either to enhance or suppress the immune responses of
other white blood cells. Other T-lymphocytes, called cytotoxic T-cells, act to kill virus-
infected cells. These different types of T-cells are distinguished from one another by the
presence of different antigenic markers on their surfaces. Specifically, helper T-cells express
20 a cell-surface glycoprotein known as CD4, and cytotoxic T-cells express a different cell-
surface glycoprotein, CD8. As T-cells mature, they express the CD2 cell-surface protein,
and, ultimately, the CD3 cell-surface protein complex.

T-cell development has been intensively studied *in vivo*. The evidence indicates that
pluripotent hematopoietic stem cells are present in populations of cells expressing CD34
25 surface molecules (CD34⁺ cells; see, for example, Terstappen et al. Blood, 79:666-677, 1992
and references cited therein). Such CD34⁺ cells represent about 1% of nucleated bone
marrow cells. CD34⁺ cells have been used to successfully repopulate the thymus of an
irradiated host with T-cells (see, for example, Exine et al. Nature 309:629-632, 1984;
Goldschneider et al. J. Exp. Med. 163:1-17, 1986; Berenson et al. J. Clin. Invest. 81:951-
30 955, 1988); and engraftment of human CD34⁺ progenitor cells into mice with genetically
determined severe combined immunodeficiency (SCID mice) has been observed to result in
development of mature T-lymphocytes in those mice (see, for example, McCune et al.
Science 241:1632-1639, 1988; Namikawa et al. J. Exp. Med. 172:1055-1063, 1990;
Berenson et al. Blood 77:1717-1722, 1991).

35 Efforts have also been directed at developing an *in vitro* T-lymphopoiesis system (see,
for example, Benveniste et al. Cell. Immunol. 127:92-104; Peault et al. J. Exp. Med.
174:1283-1286, 1991; Toki et al. Proc. natl. Acad. Sci. USA 88:7548-7551, 1991;
Tjonnfjord et al. J. Exp. Med. 177:1531-1539; Hurwitz, J. Immunol. 17:751-756, 1987).
Peault et al. have developed a system in which CD34⁺ cells from human fetal liver and bone

marrow are microinjected *in vitro* into HLA-mismatched fetal thymus fragments previously depleted of hematopoietic stem cells by low temperature culture. Peault et al. have not achieved complete *in vitro* T-cell development, however, as the *in vitro*-colonized thymuses were subsequently engrafted into SCID mice, and T-cell differentiation occurred *in vivo*.

5 Tjonnsfjord et al. have reported *in vitro* T-cell differentiation from adult human CD34+ bone marrow cells cultured in thymic stromal cell supernatant and in the presence or absence of recombinant murine *c-kit* ligand. Nevertheless, Tjonnsfjord et al. found that only a small fraction of their total cell culture (see Figure 3 of Tjonnsfjord et al.) developed into mature T-cells, as indicated by the presence of T-cell-specific surface markers such as CD2,
10 CD3, CD4, and CD8. Thus, there remains a need for the development of an *in vitro* T-lymphopoiesis system in which a significant fraction of the cultured cells develop into mature T-cells.

Summary of the Invention

15 The present invention provides an *in vitro* T-lymphopoiesis system in which a significant fraction of the cultured cells develop into mature T-cells. Approximately 17-74% of the cells produced in the *in vitro* T-lymphopoiesis system of the present invention express the T-cell-specific surface antigen CD2; 1.5-34% express CD3; 16-61% express CD4 and 0-15% express CD8. No expression of CD15 or CD56 (natural killer cell phenotype) was
20 detected.

The present invention therefore provides an isolated population of T-cells produced *in vitro* from precursor cells expressing CD34, in which approximately 17-74% of the T-cells in the population express CD2, approximately 1.5-34% of the T-cells in the population express CD3, and approximately 16-61% of the T-cells in the population express CD4. The
25 population of T-cells of the present invention expresses the lymphocyte-specific *RAG-2* gene. Furthermore, T-cells produced in the present *in vitro* system are sensitive to infection by the T-cell-tropic human immunodeficiency virus (HIV) strain HIV-1_{IIIB}. The T-cells of the present invention can therefore be utilized to identify drugs that affect HIV infection of developing T-cells.

30 The present invention also provides a method of producing T-cells *in vitro* comprising the steps of i) providing a population of precursor cells expressing CD34; and ii) culturing the precursor cells in the presence of interleukin-12 (IL-12). Preferably, the precursor cells are human bone marrow cells, human umbilical cord blood cells, human peripheral blood cells, or human fetal liver cells. The precursor cells can be cultured in the presence of a confluent
35 monolayer of thymic stroma and a cytokine or combination of cytokines including IL-12, preferably in combination with flk2/flt3 ligand. In most preferred embodiments of the method of the present invention, IL-12 is present at a concentration of about 10 ng/ml, and flk2/flt3 ligand is present at a concentration of about 100 ng/ml.

The *in vitro*-produced T-cells of the present invention can be used in pharmaceutical compositions, at therapeutically effective amounts. Preferably, such compositions include a pharmaceutically-acceptable carrier and/or a pharmaceutically-acceptable salt.

The *in vitro*-produced T-cells of the present invention can also be used to identify
5 drugs or factors that affect T-cell development. The invention also provides a combination including i) a population of CD34 positive mammalian bone marrow cells that is at least about 95 percent pure; and ii) interleukin-12, and preferably also including iii) flk2/flt3 ligand; iv) a population of thymic stromal cells; and/or v) a drug.

10 Brief Description of the Drawings

Figure 1 shows a schematic of a preferred embodiment of the method of the present invention.

Figure 2 illustrates the proliferative capabilities of various immunophenotypic subtypes of low density bone marrow mononuclear cells used as precursors in a preferred
15 embodiment of the *in vitro* T-lymphopoiesis system of present invention.

Figure 3 shows a typical flow cytometric analysis of cells produced in the *in vitro* T-lymphopoiesis system of the present invention.

Figure 4 presents PCR and Southern blot results demonstrating that CD34⁺ cells acquire *RAG-2* expression during culture in the *in vitro* T-lymphopoiesis system of the
20 present invention.

Figure 5 is a fluorescence photomicrograph showing T-cells of the present invention stained with HIV antiserum and anti-human FITC, 14 days after exposure to HIV-1_{IIIB}.

Figure 6 presents two graphs showing supernatant HIV p24 antigen levels from parallel cultures of stroma alone (panel A) and stroma with CD34⁺ cells (panel B) after
25 exposure to HIV-1_{IIIB} or heat-inactivated (mock) virus. The increase in p24 after 11 days in the cultures containing CD34⁺ cells, but not thymic stroma alone, is evidence of productive HIV infection of developing T-cells.

Figure 7 is a bar graph showing detection of IL-2 by ELISA in cultured cells of the present invention.

Figure 8 illustrates the acquisition of TCR expression by cultured cells. Day 21 cells
30 were stained with anti-CD4-PE and anti-TCR α/β -1-FITC prior to flow cytometric analysis (A). RNA derived from day 21 trypsinized thymic stroma alone, cultured CD34⁺ cells or control CD2⁺ cells was analyzed by RT PCT for TCR expression.

Figure 9 depicts the immunophenotype of cultivated cells. The output cell population
35 was gated using the forward and side scatter criteria shown (A). The flow cytometric analyses of cells stained with the indicated antibodies at particular time points are shown (B). Each time point required analysis of the entire cell population, therefore, the data shown are from independent cultures and do not represent the sequential sampling of a single culture.

Detailed Description of the Invention

As indicated above, the present invention provides an effective *in vitro* T-lymphopoiesis system. Essentially, CD34⁺ cells are plated onto a confluent monolayer of "feeder" cells, and are cultured in the presence of selected growth factors to induce T-cell differentiation. The preferred tissue source for preparation of the feeder cell layer is human fetal thymic tissue, although other sources, such as, for example, thymic stroma from non-human primates or other mammals, or immortalized cell lines derived from the above-mentioned sources may alternately be utilized. The feeder layer can be prepared using standard techniques known in the art (see, for example, Tjonnfjord et al. *supra*). The preferred method of preparation of the feeder layer is described below in Example 1.

CD34⁺ cells for use in the present invention can be obtained from a variety of different sources such as, for example, bone marrow, umbilical cord blood, peripheral blood stem, and fetal liver. The preferred source of CD34⁺ cells, due in part to its ready availability, is bone marrow. CD34⁺ cells utilized in the present invention can be purified using any available method such as, for example, fluorescence-activated cell sorter (FACS) analysis, immunomagnetic bead purification (or other immunoprecipitation method), or functional selection using cytokines in conjunction with anti-metabolites, and should be at least approximately 95% pure (i.e. at least approximately 95% of the cells should express CD34).

The purified CD34⁺ cells are plated on the feeder layer and cultured in the presence of a cytokine or mixture of cytokines, preferably including interleukin-12 (IL-12) and possibly also including flk2/flt3 ligand, for at least approximately two weeks, and typically less than or equal to approximately 35 days. Not all cytokine preparations are equally effective in the *in vitro* T-lymphopoiesis system of the present invention. In particular, we have found that IL-2, IL-7, and SCF are not as effective as is IL-12 in stimulating development of a significant fraction of cells into mature T-cells. Typically, cells proliferate in the *in vitro* system of the present invention approximately 20-25 fold in fourteen days of culture (see Figure 2).

After the culture period, cells produced in the *in vitro* system of the present invention can be analyzed for expression of T-cell markers and/or for T-cell activity using any available method including, for example, FACS analysis to detect T-cell surface markers such as, for example, CD2, CD3, CD4, CD7, CD8, CD25, and CD44. Typically, approximately 17-74% of the cells produced in the *in vitro* T-lymphopoiesis system of the present invention express detectable CD2, approximately 1.5-34% express detectable CD3, and approximately 16-61% express detectable CD4. Cell cultures produced by this *in vitro* system have fewer cells expressing CD8; that is, fewer than or equal to approximately 15% of the cells produced in the present *in vitro* system are CD8⁺.

Cultured cells of the present invention can also be analyzed for expression of T-cell-specific genes (such as, for example, *RAG-2* or T-cell receptor (TCR) genes) and/or for

production of T-cell specific proteins (e.g. IL-2) using known techniques of molecular biology (see, for Example, Sambrook et al. *Molecular Cloning: A Laboratory Manual* Cold Spring Harbor Press, NY, 1989, incorporated herein by reference). Acquisition of *RAG-2* expression is detectable during the culture period (see Figure 4). In addition, acquisition of
5 TCR β expression is detectable during the culture period (see Figure 8). IL-2 production is detectable in cultured cells produced in the *in vitro* T-lymphopoiesis system of the present invention (see Figure 7). Detection of IL-2 confirms that the cultured cells are mature T-cells.

10 T-cells produced in the *in vitro* system of the present invention can be tested for proliferative responses to mitogens such as PHA, PMA, anti-CD3 receptor (with IL-2), or for proliferative response to antigens, using known techniques.

Cultured cells of the present invention are susceptible to infection lymphotropic pathogens. In particular, cultures cells are susceptible to infection with a T-cell-tropic strain of the human immunodeficiency virus, HIV-1_{IIIB}. For example, after 14 days of culture on a
15 thymic stroma feeder layer, T-cells of the present invention that have been exposed to HIV-1_{IIIB} produce the HIV p24 antigen, that is detectable in the cell supernatant (see Figure 6). Cultured cells that are exposed to mock virus preparations, or to heat-inactivated virus, do not produce detectable p24. HIV-1_{IIIB}-exposed cells also stain positively for HIV using HIV-1 antisera (see Figure 5).

20 The present invention therefore provides an effective *in vitro* T-lymphopoiesis system. The *in vitro* system of the present invention can be utilized to study T-cell development, and also to identify potential drug candidates that interfere with or enhance a given step in differentiation and/or proliferation of T-cells. For example, drugs can be added to the culture system of the present invention, and the effect of those drugs on number,
25 immunophenotype, reactivity to known T-cell stimuli (such as, for example, PHA, PMA, anti-CD3 receptor plus IL-2), or infection by specific pathogens (such as, for example, HIV or HTLV) of the product T-cells can be determined by comparison with cultures to which no drugs have been added, or to which a "placebo" has been added. A "placebo" is a compound that does not significantly affect the parameter (e.g. cell number, etc.) being measured. The
30 specific nature of the "placebo" used in a given test depends on the character of the drug to which it is being compared. For example, if the drug in question is a protein, an appropriate placebo would be a denatured version of that protein. One of ordinary skill in the art could readily identify appropriate placebos for comparison with given drugs.

Exemplary drugs that can be tested in the *in vitro* T-lymphopoiesis system of the
35 present invention include any and all compounds suspected of having an effect on T-cell development and/or proliferation, e.g., cyclophilins. In fact, the *in vitro* T-lymphopoiesis system of the present invention can also be utilized as a source of compounds to be tested. That is, the present *in vitro* T-lymphopoiesis system, or components thereof, can be fractionated according to known procedures (see, for example, Huang et al. Cell 63:225-233,

1990; Zsebo et al. Cell 63:195-201, 1990, each of which is incorporated herein by reference), and individual fractions can be tested for their effects on T-cell development. Fractions having desirable activities (e.g. fractions that affect one or more aspects of T-cell development and/or proliferation) can be further fractionated according to known techniques, until individual active factors are identified. Active factors can subsequently be purified, and genes encoding protein factors can be cloned, according to techniques available in the art, such as those disclosed in Current Protocols in Molecular Biology (John Wiley & Sons) and Current Protocols in Immunology (John Wiley & Sons).

Active factors that affect T-cell development and/or proliferation can, for example, be derived from stromal components of the *in vitro* T-lymphopoiesis system of the present invention. Either non-immortalized or immortalized primary stromal cells can be utilized. Immortalization techniques (such as, for example, transduction with Simian Virus 40 (SV40) large T-antigen or middle T-antigen are known in the art (see, for example, Williams et al. Mol. Cell. Biol. 8:3864, 1988).

The culture system of the present invention has the additional advantage that drugs can be added at different times, so that the point during T-cell ontogeny that a given drug (or combination of drugs) exerts its effects can readily be determined.

Another valuable application for the present *in vitro* T-lymphopoiesis system and cells produced thereby is as a system for genetic manipulation of T-cells. Foreign genetic material can be introduced into, for example, the CD34⁺ precursor cells prior to culturing in the presence of a cytokine, using any available method of gene transfer such as, for example, transformation, transfection, infection (using, for example, retroviral vectors or adeno- or adeno-associated viruses), transduction, electroporation, etc. (see, for example, Sambrook et al. *supra*).

T-cells containing genes, such as, for example, dominant negative HIV-1 *rev* mutants or HIV-1 antisense constructs, that protect T-cells from infection can be produced using the *in vitro* system of the present invention. Other genes that can desirably be introduced into T-cells using the *in vitro* T-lymphopoiesis system of the present invention include, for example, genes encoding antibodies of known specificity; receptor ligand gene constructs, genes that enhance T-cell reactivity (such as genes encoding cytokines, cytokine receptors, or "co-stimulatory molecules such as, for example, CD28, B7, CD40, CD40 ligand, etc.). In some instances, it may be desirable to introduce genes that limit T-cell reactivity, so that the product T-cells of the present invention will have increased tolerance relative to T-cells that have not been subjected to similar genetic manipulation.

Additionally, the present *in vitro* T-lymphopoiesis system may be utilized to produce T-cells with specific reactivity. An antigen, such as a peptide, can be added to the culture system in order to induce proliferation of a subset of T-cells reactive against that antigen. In some cases, as will be apparent to one of ordinary skill in the art, it is desirable that the antigen be presented in the context of an antigen-presenting cell such as, for example, a T2

cell, an activated B cell, a macrophage, or a dendritic cell. Such reactive T-cells of the present invention could be utilized, for example, in adoptive immunotherapy applications in order to generate a targeted immunologic reaction for the treatment of disorders such as, for example, malignancies or infectious diseases.

5 In fact, T-cells produced in the *in vitro* system of the present invention, whether or not they have a specific reactivity, can be utilized in a variety of therapeutic techniques. In particular, the T-cells produced in the system of the present invention can be used in pharmaceutical compositions in, for example, adoptive immunotherapy, vaccine therapy and/or gene therapy. An exemplary pharmaceutical composition including a T-cell of the
10 present invention is a therapeutically effective amount of the T-cell, which may or may not have been transfected (or transformed or infected, etc.) with a gene sequence capable of expressing a particular gene product (see above), optionally combined with a pharmaceutically-acceptable and compatible carrier. The term "pharmaceutically-acceptable and compatible carrier" as used herein, and described more fully below, refers to (i) one or
15 more compatible filler diluents or encapsulating substances that are suitable for administration to a human or other animal, and/or (ii) a system capable of delivering the T-cell to a target. In the present invention, the term "carrier" thus denotes an organic or inorganic ingredient, natural or synthetic, with which the T-cells of the invention can be combined to facilitate administration.

20 The term "therapeutically-effective amount" refers to that amount of the present pharmaceutical compositions which produces a desired result or exerts a desired influence on the particular condition being treated. Various concentrations may be used in preparing compositions incorporating the same ingredient to provide for variations in the age of the patient to be treated, the severity of the condition, the duration of the treatment and/or the
25 mode of administration.

The term "compatible", as used herein, means that the components of the pharmaceutical compositions are capable of being co-mingled with the T-cells of the present invention, and with each other, in a manner such that there is no interaction that would substantially impair the desired pharmaceutical efficacy.

30 T-cells of the present invention may be administered *per se* (neat) or in combination with a pharmaceutically acceptable salt. Non-pharmaceutically acceptable salts may, in some instances, conveniently be used to prepare pharmaceutically acceptable salts, and are therefore not excluded from the scope of this invention. Pharmaceutically acceptable salts include, but are not limited to, those prepared from the following acids: hydrochloric,
35 hydrobromic, sulphuric, nitric, phosphoric, maleic, acetic, salicylic, p-toluene-sulfonic, tartaric, citric, methanesulphonic, formic, malonic, succinic, naphthalene-2-sulfonic, and benzenesulphonic. Also, pharmaceutically acceptable salts can be prepared as alkaline metal or alkaline earth salts, such as sodium, potassium or calcium salts of the carboxylic acid group.

The pharmaceutical compositions of the present invention are most suited for parenteral administration. The term "parenteral" includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques. Preferred compositions suitable for parenteral administration may comprise a sterile aqueous
5 preparation of the T-cells of the invention which is preferably isotonic with the blood of the recipient. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution.

The T-cells of the invention can also be conjugated to a moiety for use in, for example, vaccines. The moiety to which the T-cells are conjugated can be a protein,
10 carbohydrate, lipid, and the like. Coupling may be accomplished by any chemical reaction that will bind the T-cell and another molecule so long as the two moieties retain their respective activities. This linkage can include many chemical mechanisms, for instance covalent binding, affinity binding, intercalation, coordinate binding and complexation. The preferred binding is covalent binding. Covalent binding can be achieved either by direct
15 condensation of existing side chains or by the incorporation of external bridging molecules. Many bivalent or polyvalent linking agents are useful in coupling, for example, protein molecules to other molecules (e.g. molecules associated with the T-cells of the present invention).

For example, representative coupling agents can include organic compounds such as
20 thioesters, carbodiimides, succinimide esters, diisocyanates, glutaraldehydes, diazobenzenes and hexamethylene diamines. This listing is not intended to be exhaustive of the various classes of coupling agents known in the art but, rather, is exemplary of the more common coupling agents (see, for example, Killen et al. J. Immun. 133:1335-2549, 1984; Jansen et al. Immunol. Rev. 62:185-216, 1982; and Vitetta et al., supra).

The moiety to which T-cells may be bound can also be an adjuvant. The term
25 "adjuvant" is intended to include any substance which is incorporated into or administered simultaneously with the T-cells which potentiates an immune response in the subject. Adjuvants include aluminum compounds, e.g., gels, aluminum hydroxide and aluminum phosphate gels, and Freund's complete or incomplete adjuvant. The paraffin oil may be
30 replaced with different types of oils, e.g., squalene or peanut oil. Other materials with adjuvant properties include BCG (attenuated Mycobacterium tuberculosis plus other microbial derivatives), calcium phosphate, levamisole, isoprinosine, polyanions (e.g., poly A:U), leutinin, pertussis toxin, lipid A, saponins and peptides, e.g., muramyl dipeptide. Rare earth salts, e.g., of lanthanum and cerium, may also be used as adjuvants. The amount of
35 adjuvant required depends upon the subject and the particular therapeutic used and can be readily determined by one skilled in the art without undue experimentation. An exemplary pharmaceutical composition of the present invention therefore can comprise a therapeutically effective amount of the T-cells of the invention, either alone or in combination with an immunoadjuvant and/or carrier.

Because the cell cultured in the present *in vitro* system are susceptible to infection by HIV-1, this system can be used to identify drugs that inhibit HIV-1 infection and/or replication, particularly in developing T-cells. The present *in vitro* system can also be used to identify drugs that inhibit development and/or proliferation of HIV-infected cells. For example, standard virus infection techniques can be employed to infect cells in the *in vitro* cultures of the present invention with laboratory strains of HIV-1 (such as, for example, HIV-1_{IIIB}) or with primary patient isolates from HIV-1-infected individuals. Potential drugs can be added to the cultures at different points during infection and the effects of those drugs on viral infection can be assayed using standard HIV-1 detection systems (such as, for example, assays for HIV-1 p24 antigen, reverse transcriptase assays, immunofluorescence assays, or PCR-based assays) and comparing the results found with infected cultures that are not treated with drugs, or that are treated with a placebo as discussed above. For examples of some potential drugs that could be analyzed using the present system, and of methods for assaying the effects of those drugs on HIV infection, see, for example, Tsubota et al. J. Clin. Invest. 86:1684-1689, 1990; Gao et al. Proc. Natl. Acad. Sci. USA 90:8925-8928, 1993; and Meyerhans et al. J. Virol. 68:535-540, 1994.

This invention is further illustrated by the following examples which in no way should be construed as being further limiting. The contents of all cited references (including literature references, issued patents, published patent applications, and co-pending patent applications) cited throughout this application are hereby expressly incorporated by reference.

EXAMPLES

EXAMPLE 1: PREPARATION AND ANALYSIS OF T-CELLS PRODUCED *IN VITRO*

Thymic stromal culture

Thymus glands were removed from electively aborted human fetuses of 18-24 weeks gestation after informed maternal consent (Anatomic Gift Foundation, Laurel, MD) and transported at room temperature in culture medium. Thymic tissue was minced into small fragments using scalpel blades and then passed through a tissue sieve to obtain a single cell suspension. Cells were washed twice in phosphate buffered saline (PBS) and resuspended in Iscove's modified Dulbecco's medium (IMDM) (Mediatech, Washington, DC) containing 20% FCS (Sigma, St. Louis, MO), glutamine, penicillin and streptomycin at a density of 2×10^6 /ml. The cell suspension was plated into 24-well tissue culture plates, 1 ml per well, and cultured for 10-14 days at 37°C in a humidified incubator containing 5% CO₂ to allow a

confluent stromal monolayer to form. Thymic cells not used immediately were cryopreserved in 10% DMSO and washed in medium after thawing prior to plating out as above.

5 CD34⁺ cell purification from adult bone marrow and fetal liver

Bone marrow samples were obtained with informed consent from healthy adult volunteers by iliac crest aspiration. Fetal livers (18-24 weeks gestation) were obtained as described above for the thymus glands and single cell suspensions were prepared in the same manner. Mononuclear cells (MNC) were separated from the marrow or liver samples by
10 centrifugation over Ficoll-Paque (Pharmacia Biotechnology, Uppsala, Sweden), washed twice and suspended in IMDM. Bone marrow MNC were cultured overnight to remove adherent cells prior to further purification.

In order to isolate the CD34⁺ fraction, the cells were first incubated for one hour to 4°C with mouse anti-human CD34 IgG1 monoclonal antibody (Amac Inc., Westbrook, ME),
15 washed twice in IMDM and then incubated with Dynabeads M-450 coated with anti-mouse IgG (DynaI Inc., Great Neck, NY) for one hour at 4°C with gentle agitation. The selected cells were washed four times using a magnet and their purity checked by microscopy. In some experiments, bone marrow MNC were also depleted of cells of T lineage by negative beads selection (via two rounds of negative selection with Dynabeads M-450 coated with
20 mouse anti-CD2 monoclonal IgG1 antibody (DynaI Inc.), or via use of anti-CD34 DETACHABEADSTM (DynaI, Inc.)), or by cell sorting. CD2⁺CD34⁻ and CD2⁺CD34⁺ fractions were purified by FACS for use in control experiments.

25 Purification of marrow cell fractions by fluorescence-activated cell sorter (FACS) analysis

Bone marrow MNC were stained with FITC-conjugated anti-CD34 and PE-conjugated anti-CD2 antibodies, as described below. Stained cells were sorted using a FACS Vantage cell sorter (Becton Dickinson, San Jose, CA); two color fluorescence was quantitated with log amplification in the FL1 and FL2 channels and the CD34⁺ CD2⁺ and
30 CD34⁺ CD2⁻ fractions collected separately.

CD34⁺ cultures

Between 5×10^4 - 10^5 CD34⁺ per well, purified as described above, were plated on top of the thymic stromal monolayers which were first washed extensively to remove all non-
35 adherent cells, and, in some experiments, were irradiated with 15 Gy. The cells were then cultured for a further 2-4 weeks in IMDM/20% FCS with and without added cytokines, as described below. Control cultures were performed using either bone marrow stromal monolayers or empty wells with no stroma instead of the thymic stroma. Cultures were fed

twice weekly by carefully removing most of the medium from each well and replacing it with 1 ml fresh medium and appropriate cytokines.

Cytokines

- 5 The following human cytokines were added in combination to achieve optimal conditions for promoting T lineage differentiation: - 5%, Interleukin-12 (IL-12), 10 ng/ml (R&D Systems, Minneapolis, MN) and flk2/flt3 ligand, 100/ng (a kind gift from Dr. S. Lyman, Immunex, Seattle, WA).

10 Immunophenotyping of cultured cells by flow cytometry

- Cells were harvested from the culture plates at various times between 2-4 weeks after plating by gentle aspiration, so as to leave the stromal cells behind. The cells were counted and their viability assessed by trypan blue staining. Aliquots of between 2.5×10^5 cells were washed twice in PBS and stained with directly conjugated (FITC or PE) mouse monoclonal
15 antibodies to human surface antigens. The antibodies used were as follows: -FITC-anti-CD34 (Amac Inc.), FITC-anti-CD2 and anti-CD7, PE-anti-CD2, anti-CD3, anti-CD4, anti-CD7 and anti-CD8 (Exalpha Corp., Boston, MA), FITC-anti-TCR α/β 1 (Becton Dickinson), and FITC-anti-CD56 + anti-CD16. FITC- and PE- conjugated mouse isotype control antibodies were used for each culture. Staining was performed according to the
20 manufacturer's instructions and cells were fixed in 1% paraformaldehyde prior to analysis by flow cytometry. Flow cytometric analysis was performed on 10,000 cells from each sample using a FACScan cytometer (Becton Dickinson). Dual color immunofluorescence was quantitated using log amplification in the FL1 and FL2 channels and analyzed using LYSYS II (Becton Dickinson) software for the Hewlett Packard 9000 series computer.

- 25 Figure 3 shows a typical flow cytometric analysis of a CD2-depleted, CD34⁺ cell culture of the present invention that has been supplemented with flk2/flt3 ligand and IL-12, harvested at 21 days. There was considerable variability between cultures, but the combination of flk2/flt3 ligand and IL-12 consistently yielded the highest fraction of cells expressing T cell markers. Culture of purified CD34⁺ cells for 14-35 days with these
30 cytokines resulted in 17-74% CD2, 1.5-34% CD3 and 16-61% CD4 expression. Expression of CD7 was lower than that of CD2 in most experiments. CD8 expression was low ($\leq 15\%$) in most of the cultures, and there was no detectable expression of CD16 or CD56 (natural killer cell markers).

- 35 Figure 9 shows other typical flow cytometric analyses of CD34⁺, CD2⁻ bone marrow cells cultured in the described system for the internal indicated.

PMA/PHA-stimulated IL-2 production by cultured cells

In order to confirm the presence of relatively mature T cells in the harvested cells, aliquots of 10^5 cells in 200 μ l IMDM/20% FCS were cultured with and without the addition

of phytohemagglutinin (PHA), 3 µg/ml, and phorbol myristic acid (PMA), 50 ng/ml (both Sigma). The CD2⁺ fraction of bone marrow MNC, separated as described above, was used as a positive control. The culture supernatants were harvested after 48h and stored at -20°C before assaying the interleukin-2 (IL-2) concentration by ELISA (R & D Systems). Low, but
5 detectable, levels of IL-2 were observed in the culture supernatants.

Reverse-transcription-coupled polymerase chain reaction (RT-PCR) for *RAG-2* and *IL-2* gene expression

Total RNA was prepared from between 10³-10⁴ sorted or cultured cells by
10 guanidinium isothiocyanate extraction and pelleted by ultracentrifugation through cesium chloride. cDNA was prepared by reverse transcription using random primers and Moloney reverse transcriptase (Gibco-BRL), and was amplified using primers specific for a 415 basepair (bp) region of the human *RAG-2* gene, which is expressed transiently only by cells
15 undergoing lymphocyte differentiation. PCR products were analyzed by 2% agarose gel electrophoresis and photographed under UV light after ethidium bromide staining (see Figure 4). Specificity was confirmed by Southern blot hybridization with a ³²P-labelled 32 base probe, followed by autoradiography.

EXAMPLE 2: INFECTION OF *IN VITRO*-PRODUCED T-CELLS WITH HIV-2_{IIIB}

20

Cell-free HIV-1_{IIIB} or mock virus preparations were added to *in vitro* T-cell cultures to determine susceptibility to HIV-1 infection. Supernatants were collected at intervals of 3-4 days and HIV p24 antigen levels were measured by ELISA.

25 Equivalents

It should be understood that the preceding is merely a detailed description of certain preferred embodiments of the present invention. It therefore should be apparent to those skilled in the art that various modifications and equivalents can be made without departing from the spirit or scope of the invention.

30

What is claimed is:

1. A population of T-cells produced *in vitro* from precursor cells expressing CD34, wherein approximately 17-74% of the T-cells in said population express CD2, approximately 1.5-34% of the T-cells in said population express CD3, approximately 16-61% of the T-cells in said population express CD4, and approximately 0-15% of the T-cells in said population express CD8.

2. The population of T-cells of claim 1 wherein at least one cell in said population contains foreign genetic material.

3. The population of T-cells of claim 1 wherein at least one cell in said population is infected with HIV.

4. A method of producing T-cells *in vitro* comprising:
providing a population of precursor cells expressing CD34; and
culturing said precursor cells in the presence of interleukin-12 (IL-12).

5. The method of claim 4 wherein the step of providing a population of precursor cells comprises providing a population of human cells selected from the group consisting of: bone marrow cells, fetal liver cells, umbilical cord blood cells or peripheral blood cells.

6. The method of claim 4, wherein the step of providing comprises providing a population of precursor cells that is at least about 95 percent pure.

7. The method of claim 6, wherein the step of providing a 95 percent pure population of precursor cells comprises:
providing a cell population including said precursor cells;
removing adherent cells from said cell population;
isolating said 95 percent pure population of precursor cells from said cell population by immunomagnetic bead isolation or fluorescence activated cell sorting (FACS).

8. The method of claim 7, wherein the step of providing a 95 percent pure population of precursor cells further comprises:
depleting said immunomagnetic bend or FACS isolated population of cells of T-cell lineage by a method selected from the group consisting of negative bead selection and cell sorting.

9. The method of claim 4, further comprising the step of providing a confluent monolayer of thymic stroma to said precursor cells, and wherein the step of culturing

comprises culturing said precursor cells in the presence of interleukin-12 and in contact with said confluent monolayer.

5 10. The method of claim 9, wherein the step of providing a confluent monolayer comprises:

 providing human fetal thymic tissue;
 preparing said confluent monolayer from cells from said tissue.

10 11. The method of claim 4, wherein the step of culturing comprises culturing said precursor cells in the presence of interleukin-12 and flk2/flt3 ligand.

15 12. The method of claim 11, wherein the step of culturing comprises culturing said precursor cells in the presence of interleukin-12 at about 10 ng/ml and flk2/flt3 ligand at about 100 ng/ml.

 13. A pharmaceutical composition comprising a therapeutically effective amount of T-cells of claim 1.

20 14. The composition of claim 13, further comprising a pharmaceutically-acceptable carrier.

 15. The composition of claim 14, further comprising a pharmaceutically-acceptable salt.

25 16. In combination, a population of CD34 positive mammalian bone marrow cells that is at least about 95 percent pure and interleukin-12.

 17. The combination of claim 16 further comprising flk2/flt3 ligand.

30 18. The combination of claim 16, further comprising a population of thymic stromal cells.

 19. The combination of claim 18, further comprising a population of thymic stromal cells.

35 20. The combination of claim 19, further comprising a drug.

 21. The combination of claim 20, wherein at least one cell of said population is infected with HIV.

22. Active factors identified by the method of claim 4 derived from a population of T-cells or thymic stromal cells.

5 23. A method for identifying active factors affecting T-cell development and/or proliferation, said method comprising:

producing T-cells *in vitro* from a population of precursor cells expressing CD34, fractionating components of at least one of

- 10 i) said T-cells, or products thereof,
ii) thymic stromal cells, or products thereof,

to produce a candidate factor,

applying said candidate factor to a population of precursor cells expressing CD34 to identify an active factor affecting T-cell development and/or proliferation.

15 24. A method according to claim 23, wherein said producing step includes the step of culturing said precursor cells in the presence of interleukin-12 (IL-12).

25 25. Active factors identified by the method of producing T-cells *in vitro* from a population of precursor cells expressing CD34, fractionating components of at least one of

- 20 i) said T-cells, or products thereof,
ii) thymic stromal cells, or products thereof,

to produce a candidate factor,

25 applying said candidate factor to a population of precursor cells expressing CD34 to identify an active factor affecting T-cell development and/or proliferation.


30 26. Active factors identified by the method of producing T-cells *in vitro* from a population of precursor cells expressing CD34, culturing said precursor cells in the presence of interleukin-12 (IL-12); fractionating components of at least one of


- i) said T-cells, or products thereof,
ii) thymic stromal cells, or products thereof,

to produce a candidate factor,

35 applying said candidate factor to a population of precursor cells expressing CD34 to identify an active factor affecting T-cell development and/or proliferation.

1 / 9

*FIG. 1*

*FIG. 2*






Cell Phenotype
Day 21

FIG. 3



**FIG. 4****FIG. 5**

RECTIFIED SHEET (RULE 91)

5 / 9

**FIG. 6A****FIG. 6B**

RECTIFIED SHEET (RULE 91)

**FIG. 7****FIG. 8A**




FIG. 8B



*FIG. 9A*

FIG. 9B



RECTIFIED SHEET (RULE 91)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/15307

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : A61K 45/05; A01N 63/00; C12N 5/00

US CL : 424/85.2, 93.71; 435/240.2

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/85.2, 93.71; 435/240.2

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, MEDLINE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, E	US, A, 5,476,780 (WATANABE) 19 December 1995, see entire document.	1-3, 13-15
X	Journal of Experimental Medicine, Volume 177, issued June 1993, Tjonnfjord et al., "T Lymphocyte Differentiation In Vitro from Adult Human Prethymic CD34 ⁺ Bone Marrow Cells", pages 1531-1539, see abstract.	1
--		--
Y		2, 3, 13-15
X	Cellular Immunology, Volume 127, issued 1990, Benveniste et al., "Development of T Cells <i>in vitro</i> from Precursors in Mouse Bone Marrow", pages 92-104, see abstract.	1
--		--
Y		2, 3, 13-15



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	
A document defining the general state of the art which is not considered to be of particular relevance	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
E earlier document published on or after the international filing date	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
O document referring to an oral disclosure, use, exhibition or other means	*A* document member of the same patent family
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

06 MARCH 1996

Date of mailing of the international search report

25 MAR 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

Sandra Saucier

Telephone No. (703) 308-6196

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/15307

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X -- Y	Journal of Immunology, Volume 147, Number 3, issued 01 August 1991, Gately et al., "Regulation of human lymphocyte proliferation by a heterodimeric cytokine, IL-12 (cytotoxic lymphocyte maturation factor)", pages 874-882, see abstract.	1 -- 2, 3, 13-15
X -- Y	Journal of Immunology, Volume 149, Number 11, issued 01 December 1992, Perussia et al., "Natural killer cell stimulatory factor or IL-12 has differential effects on the proliferation of TCR- $\alpha\beta^+$, TCR- $\gamma\delta^+$ T lymphocytes and NK cells", pages 3495-3502, see abstract.	1 -- 2, 3, 13-15
X -- Y	Journal of Immunology, Volume 151, Number 6, issued 15 September 1993, Cesano et al., "Cellular and molecular mechanisms of activation of MHC nonrestricted cytotoxic cells by IL-12", pages 2943-2957, see abstract.	1 -- 2, 3, 13-15
X	Science, Volume 262, Number 5140, issued 10 December 1993, Clerici et al., "Restoration of HIV-specific cell-mediated immune responses by interleukin-12 in vitro", pages 1721-1724, see abstract.	1-3, 13-15

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/15307

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1-3, 13-15

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1.

Group I, claims 1-3, 13-15, drawn to a first composition comprising a population of T cells wherein 17-74% express CD2, 1.5-34% express CD3, 16-61% express CD4 and 0-15% express CD8.

Group II, claims 4-12, drawn to a first method of making T-cells in vitro by culturing precursor cells with IL-12.

Group III, claims 16-21, drawn to a second composition comprising CD34 positive bone marrow cells at least 95% pure and IL-12.

Group IV, claims 22 and 25, drawn to a third composition comprising active factors.

Group V, claims 23-24, drawn to a second method of identifying active T-cell factors.

The inventions listed as Groups I-IV do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: Groups I, III and IV are drawn to distinct compositions which require distinct components and, therefore, are not related because they lack the same special technical feature. For example, the T-cells of Group I are required to express specific amounts of specific cell surface antigens which are not required for the compositions of groups III or IV. Group III requires IL-2 as a component of the composition which is not required by Groups I or IV. Group IV is an active factor which is not required by the compositions of Group I or III.

Groups II and V are two distinct methods. Group II is drawn to a method of making T-cells with certain characteristics, while Group V is a method for identifying active factors. Hence the two methods have distinct end points and have distinct steps in each method.

The methods of Groups II or V are not related by a special technical feature to the compositions of Groups I, III or IV because the methods of Groups II or V are not methods of either using or making any of the compositions of Groups I, III or IV as claimed.

PCT Rule 13.2 defines the permitted categories as

First appearing product

First appearing process of making the First product

First appearing method of using the First product

OR

First appearing process

First appearing apparatus for performing First process

OR

First appearing product

First appearing process of making First product

First appearing apparatus for performing First process.

Because the inventions as claimed do not match one of the permitted categories, and they are not so linked by a special technical feature within the meaning of PCT Rule 13.2 so as to form a single inventive concept, the holding of Lack of Unity as indicated above is deemed proper.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☒ **FADED TEXT OR DRAWING**
- ☒ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☒ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.